

# West-wide Jumpstart Air Quality Modeling Study

## Final Project Report and Modeling Results



September 23, 2014

Tom Moore

WRAP Air Quality Program Manager

Technical Project Team

ENVIRON, Alpine Geophysics, Univ. of North Carolina

Funding from State of NM, BP, and National BLM Air Program

Oversight by western states, local air agencies, federal land managers, EPA regional and national offices



# EPA national Ozone Standard

- Measured at ground station sites, highest 8-hour average each day
- 4<sup>th</sup> highest values each year are averaged over 3-year periods to determine compliance (e.g., 2007-09, 2008-10)
  - Statistic is called a “Design Value” for that site for that time period
- Current Ozone health standard level is 75 ppb
- EPA (re)considering revised Ozone health standard in a range of 60 to 70 ppb
- EPA also considering a secondary Ozone standard for ecosystem protection
  - Growing season / daylight hours-weighted cumulative metric

# What are (some of) the sources and control issues in the West related to new Ozone standard(s)?


- Urban and rural reactivity
- Transport and formation – how much / how important?
- Public lands with large biogenic emissions and fire activity
  - How to characterize for effects of drought and climate variation ?
- Federal and state mobile fuel and tailpipe controls
- Upstream Gas NSPS rules in place in 2015
  - Industry practices changing rapidly, e.g., green completions
- Point sources (dominated by EGUs for  $\text{SO}_2$ ,  $\text{NO}_x$  )
  - Significant  $\text{NO}_x$  BART by ~2018
  - Less coal-fired electricity supply due to climate change rule?
  - 17+ million acres of public lands leased in last 5 years for O&G exploration and production

# Introduction

- West-wide Jump-start Air Quality Modeling Study (WestJumpAQMS) was initiated in late 2010 to:
  - Develop the next generation of regional air quality modeling databases for ozone, PM<sub>2.5</sub>, visibility and deposition planning in the western U.S.
  - Provide information on the role of interstate and international transport to ozone and PM<sub>2.5</sub> under current and potential future NAAQS
  - Assess contributions of major source categories (e.g., point, O&G, mobile, et cetera) to air quality in the West
  - Provide detailed information to the community

# Overview of Approach

- 2008 Modeling Database
  - 36 km CONUS
  - 12 km WESTUS
  - 4 km IMWD
- WRF meteorological; CAMx photochemical; SMOKE emissions models
- 2008 WRAP Phase III O&G emissions
- 2008 NEI emissions
- Model Evaluation
- Sensitivity Tests



- State-Specific and Source Category-Specific Ozone and PM<sub>2.5</sub> Source Apportionment Modeling

# WestJumpAQMS Products

- Final Report
  - 15 Electronic Appendices
  - Response-to-Comments
- Ammonia Emissions Recommendations Memo
- Modeling Protocol
  - Response-to-Comments
- WRF Application/Evaluation Report
  - Evaluation down to individual monitoring site
  - Response-to-Comments
- 16 Technical Memorandums on Emissions
  1. Point Sources
  2. Area + Non-Road
  3. On-Road Mobile
  - 4a-e. Oil and Gas (5 geographic areas)
  5. Fire (WF, Rx & Ag)
  6. Fugitive Dust
  7. Off-Shore Shipping
  8. Ammonia
  9. Biogenic
  - 11 Mexico/Canada
  12. Sea Salt and Lightning
  13. Emissions Modeling Parameters

**All information on WestJumpAQMS web page:**

**<http://www.wrapair2.org/WestJumpAQMS.aspx>**

# Ozone, PM, Deposition, and Visibility Source Apportionment Resources from WestJumpAQMS

**WestJumpAQMS - Reports**

West-Wide Jump-Start Air Quality Modeling Study (WestJumpAQMS) – Final Report ([PDF](#) 15MB), September 30, 2013

- Response-to-Comments for Draft Final Report ([PDF](#) 1MB), September 30, 2013

List of Appendices and directions for use ([PDF](#))

Appendix A: CSAPR-Type Analysis for 2008 Upwind State Highest Contribution to Average and Maximum Ozone Design Values at any Monitoring Site in up to 5 Downwind States using MATS ([XLSX](#) 1MB)

Appendix B: State Contributions to Daily Maximum 8-Hour Ozone Concentrations on 10 Highest Modeled Ozone Days in 2008 by Monitoring Site ([XLSX](#) 19MB)

Appendix C: Spatial Maps of State-Specific Anthropogenic Emissions Contributions to Highest and Fourth Highest Modeled Daily Maximum 8-Hour Ozone Concentrations during 2008 Greater than 76 (current NAAQS), 70, 65, 60 (potential future NAAQS) and 0 (maximum contribution) ppb across the 12 km WESTUS and 36 km CONUS Domains ([ZIP](#) 37MB) (**corrected files posted February 7, 2014**)

Appendix D: CSAPR-Type Analysis for 2008 Upwind State Highest Contribution to Annual PM<sub>2.5</sub> Design Values in up to Five Downwind States using MATS ([XLSX](#) 12MB)

Appendix E: State Contributions to Modeled Annual PM<sub>2.5</sub> Concentrations in 2008 by Monitoring Site ([XLSX](#) 23MB)

Appendix F: CSAPR-Type Analysis for 2008 Upwind State Highest Contribution to 24-Hour PM<sub>2.5</sub> Design Values in up to Five Downwind States using MATS ([XLSX](#) 12MB)

Appendix G: Spatial Maps of Modeled State-Specific Anthropogenic Emissions Contributions to Highest (1stmax) and Eighth (8thmax) Highest 24-Hour PM<sub>2.5</sub> Concentrations during 2008 greater than 35 (current NAAQS), 30, 25, 20 and 0 (maximum contribution)  $\mu\text{g}/\text{m}^3$  ([ZIP](#) 13MB) (**corrected files posted February 7, 2014**)

Appendix H: Source Category-Specific Contributions to 8-Hour Ozone Design Values at Monitoring Sites in the 4 km Detailed Source Apportionment Domain (DSAD) using MATS and Maximum Contribution to 8-Hour Ozone Design Values in Each DSAD State (CO, NM, UT and WY) due to Major Source Categories using MATS ([XLSX](#) 1MB)

Appendix I: Source Category-Specific Contributions to Ten Highest Modeled Daily Maximum 8-Hour Ozone Concentrations at Monitoring Sites in the 4 km Detailed Source Apportionment Domain (DSAD) ([XLSX](#) 2MB)

Appendix J: Source Category-Specific Contributions to Annual PM<sub>2.5</sub> Design Values at Monitoring Sites in the 12 km WESTUS Domain using MATS ([XLSX](#) 2MB)

Appendix K: Source Category-Specific Contributions to Modeled Annual PM<sub>2.5</sub> Concentrations ( $\mu\text{g}/\text{m}^3$ ) at Monitoring Sites in the 12 km WESTUS Domain ([XLSX](#) 4MB)

Appendix L: Source Category-Specific Contributions to 24-Hour PM<sub>2.5</sub> Design Values at Monitoring Sites in the 12 km WESTUS Domain using MATS ([XLSX](#) 2MB)


Appendix M: Source Category-Specific Contributions to Ten Highest Modeled 24-Hour PM<sub>2.5</sub> Concentrations ( $\mu\text{g}/\text{m}^3$ ) at Monitoring Sites in the 12 km WESTUS Domain ([XLSX](#) 10MB)

Appendix N: Annual Sulfur and Nitrogen Wet and Dry Deposition at IMPROVE Monitors by Species ([XLSX](#) 1MB)

Appendix O: Western State-Specific Modeled Contributions to Visibility Impairment at IMPROVE Monitoring Sites for Modeled Worst (W20) and Best (B20) 20% Days during 2008 ([ZIP](#) 46MB)




# Ozone Attainment Test Software – Unmonitored Area Analysis with Design Value (2006-2010) $\geq 76$ ppb




○ Min(210,3) = 76.00, ◇ Max(45,67) = 113.30

# Ozone Attainment Test Software – Unmonitored Area Analysis with Design Value (2006-2010) $\geq 70$ ppb




# Attainment Test Software – Unmonitored Area Analysis with Design Value (2006-2010) $\geq 65$ ppb




○ Min(177,1) = 65.00, ◇ Max(45,67) = 113.30

# Ozone Attainment Test Software – Unmonitored Area Analysis with Design Value (2006-2010) $\geq 60$ ppb








*"They have very strict anti-pollution laws in this state."*

# State-Specific Ozone Source Apportionment

- Purpose: To provide information on the role of ozone transport to exceedances of the current and potential future ozone NAAQS in the western U.S.
- Approach: Analyze ozone apportionment several ways:
  1. Upwind state contribution to downwind state nonattainment using Cross State Air Pollution Rule (CSAPR-type) approach
    - Use EPA method for projecting ozone Design Values (RRFs)
  2. State contributions to modeled high ozone DMAX8 ozone at monitors in 12 km WESTUS domain
    - Spatial extent of modeled state contributions to 1stmax and 4thmax DMAX8 ozone greater than current and potential future NAAQS
    - Source category analysis (Natural, Fires & Anthropogenic)
  3. Detailed Source Category-Specific Source Apportionment
    - 6 key source categories across 4 states in intermountain West
    - 2-way nesting between model domains

# State-Specific Ozone Source Apportionment

- 2008 36/12 km Base
- 17 Western States
  - Plus EasternUS, Can, Mex & Off-Shore
- 5 Source Categories
  - Natural  
(Biogenics+Lightning+WBDust+SeaSalt)
  - WF, Rx and Ag Fires
  - Anthropogenic
- 107 Source Groups ( $21 \times 5 + 2$ )
  - 4 Extra Species for each Group
    - 428 additional species
    - Standard Model = 70 species
    - Computationally Demanding





# CSAPR-Type Analysis for current (76 ppb) and potential future (70 and 65 ppb) NAAQS levels

- CSAPR looked at contributions to:
  - Average Design Value = Average of DVs from 2006-2010
  - Max Design Values = Max DVs from 2006-2010
- 136 ozone monitors in 12 km WESTUS domain with Average Design Value exceeding 76 ppb NAAQS
  - 86 sites (63%) in California
- For 17 upwind western states examine 2008 contribution to DMAX8 ozone Design Value in downwind states
  - CSAPR used a 1% NAAQS significance threshold ( $\geq 0.76$  ppb)
- This analysis is for 2008 and is not a regulatory analysis that would have to examine a future year




# Nevada CSAPR-Type Ozone Analysis for current 75 ppb NAAQS (from WestJumpAQMS Appendix A)




# Nevada CSAPR-Type Ozone Analysis for potential 70 and 65 ppb NAAQS (from WestJumpAQMS Appendix A)

NEVADA's Contributions to Top 5 States Surrounding NEVADA, where DV is at or above 70.0 ppb Threshold




NEVADA's Contributions to Top 5 States Surrounding NEVADA, where DV is at or above 65.0 ppb Threshold




# Nevada CSAPR-Type Ozone Analysis for potential 60 ppb NAAQS (from WestJumpAQMS Appendix A)

States' Contributions to Top 5 States Surrounding NEVADA, where DV is at or above 60.0 ppb Threshold




NEVADA's Contributions to Top 5 States Surrounding NEVADA, where DV is at or above 60.0 ppb Threshold




# Nevada CSAPR-Type Ozone Contribution Analysis in “states near NV” at potential 65 ppb NAAQS (from WestJumpAQMS Appendix A)


NEVADA's Contributions to Top 5 States Surrounding CALIFORNIA, where DV is at or above 65.0 ppb Threshold




NEVADA's Contributions to Top 5 States Surrounding OREGON, where DV is at or above 65.0 ppb Threshold



NEVADA's Contributions to Top 5 States Surrounding UTAH, where DV is at or above 65.0 ppb Threshold




NEVADA's Contributions to Top 5 States Surrounding ARIZONA, where DV is at or above 65.0 ppb Threshold



# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**Highest Modeled DMAX8 Day at “Reno3” on State Street, Reno**

**High Day Contributions to MDA8 Ozone [ppb]**



**Site: NV\_Washoe0016**

**Rank: 1 - 10 Jul, 2008**

**Total Ozone = 103.4 ppb**


**BC Ozone = 44.2 ppb (42.8%)**

AZ	AZ (0.01 ppb, 0.01%)
CA	CA (5.09 ppb, 4.92%)
CO	CO (0.00 ppb, 0.00%)
KS	KS (0.00 ppb, 0.00%)
ID	ID (0.01 ppb, 0.01%)
MT	MT (0.00 ppb, 0.00%)
OK	OK (0.00 ppb, 0.00%)
OR	OR (1.43 ppb, 1.38%)
WA	WA (0.54 ppb, 0.52%)
WY	WY (0.00 ppb, 0.00%)
ND	NE (0.00 ppb, 0.00%)
SD	NV (0.16 ppb, 0.15%)
NE	UT (0.01 ppb, 0.01%)
NV	TX (0.00 ppb, 0.00%)
UT	NM (0.00 ppb, 0.00%)
TX	Eastern (0.00 ppb, 0.00%)
NM	Canada (0.02 ppb, 0.02%)
Eastern	Mexico (0.00 ppb, 0.00%)
Canada	Ocean (0.04 ppb, 0.04%)
Mexico	Natural (2.25 ppb, 2.17%)
Ocean	Wild Fire (49.60 ppb, 47.3%)
Natural	Rx Fire (0.00 ppb, 0.00%)
Wild Fire	AG Fire (0.00 ppb, 0.00%)
Rx Fire	
AG Fire	

# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**4<sup>th</sup> Highest Modeled DMAX8 Day at “Reno3” on State Street, Reno**

**High Day Contributions to MDA8 Ozone [ppb]**



**Site: NV\_Washoe0016**

**Rank: 4 - 24 Jun, 2008**

**Total Ozone = 74.5 ppb**


**BC Ozone = 33.5 ppb (44.9%)**

AZ	AZ (0.00 ppb, 0.00%)
CA	CA (17.13 ppb, 23.00%)
CO	CO (0.00 ppb, 0.00%)
KS	KS (0.00 ppb, 0.00%)
ID	ID (0.00 ppb, 0.00%)
MT	MT (0.00 ppb, 0.00%)
OK	OK (0.00 ppb, 0.00%)
OR	OR (0.26 ppb, 0.35%)
WA	WA (0.00 ppb, 0.00%)
WY	WY (0.00 ppb, 0.00%)
ND	ND (0.00 ppb, 0.00%)
SD	SD (0.00 ppb, 0.00%)
NE	NE (0.00 ppb, 0.00%)
NV	NV (0.00 ppb, 0.00%)
UT	UT (0.00 ppb, 0.00%)
TX	TX (0.00 ppb, 0.00%)
NM	NM (0.00 ppb, 0.00%)
Eastern	Eastern (0.00 ppb, 0.00%)
Canada	Canada (0.00 ppb, 0.00%)
Mexico	Mexico (0.00 ppb, 0.00%)
Ocean	Ocean (0.28 ppb, 0.38%)
Natural	Natural (0.97 ppb, 1.31%)
Wild Fire	Wild Fire (21.99 ppb, 29.40%)
Rx Fire	Rx Fire (0.36 ppb, 0.49%)
AG Fire	AG Fire (0.01 ppb, 0.01%)

# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**10<sup>th</sup> Highest Modeled DMAX8 Day at “Reno3” on State Street, Reno**

## High Day Contributions to MDA8 Ozone [ppb]



Site: NV\_Washoe0016

Rank: 10 - 03 May, 2008


Total Ozone = 70.3 ppb

BC Ozone = 61.4 ppb (87.4%)

AZ	AZ (0.00 ppb, 0.00%)
CA	CA (3.86 ppb, 5.49%)
CO	CO (0.00 ppb, 0.00%)
KS	KS (0.00 ppb, 0.00%)
ID	ID (0.00 ppb, 0.00%)
MT	MT (0.00 ppb, 0.00%)
OK	OK (0.00 ppb, 0.00%)
OR	OR (0.17 ppb, 0.25%)
WA	WA (0.03 ppb, 0.04%)
WY	WY (0.00 ppb, 0.00%)
ND	ND (0.00 ppb, 0.00%)
SD	SD (0.00 ppb, 0.00%)
NE	NE (0.00 ppb, 0.00%)
NV	NV (4.00 ppb, 5.69%)
UT	UT (0.00 ppb, 0.00%)
TX	TX (0.00 ppb, 0.00%)
NM	NM (0.00 ppb, 0.00%)
Eastern	Eastern (0.00 ppb, 0.00%)
Canada	Canada (0.01 ppb, 0.01%)
Mexico	Mexico (0.00 ppb, 0.00%)
Ocean	Ocean (0.07 ppb, 0.11%)
Natural	Natural (0.63 ppb, 0.89%)
Wild Fire	Wild Fire (0.10 ppb, 0.14%)
Rx Fire	Rx Fire (0.01 ppb, 0.02%)
AG Fire	AG Fire (0.00 ppb, 0.00%)

# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)


**Highest Modeled DMAX8 Day at 4<sup>th</sup> Street site, Sparks**






# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**4<sup>th</sup> Highest Modeled DMAX8 Day at 4<sup>th</sup> Street site, Sparks**




# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**10<sup>th</sup> Highest Modeled DMAX8 Day at 4<sup>th</sup> Street site, Sparks**




# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

## Highest Modeled DMAX8 Day at Great Basin NP, White Pine County




# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

## 4th Highest Modeled DMAX8 Day Great Basin NP, White Pine County



# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

## 10th Highest Modeled DMAX8 Day Great Basin NP, White Pine County






# Clark County monitoring sites



# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)


**Highest** Modeled DMAX8 Day at Jean background site on SR 161, southwest of Las Vegas





# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)


**4<sup>th</sup> Highest** Modeled DMAX8 Day at Jean background site on SR 161, southwest of Las Vegas






# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**10<sup>th</sup> Highest Modeled DMAX8 Day at Jean background site on SR 161, southwest of Las Vegas**




# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**Highest** Modeled DMAX8 Day at Joe Neal site, Azure Street, northwest Las Vegas




# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**4<sup>th</sup> Highest** Modeled DMAX8 Day at Joe Neal site, Azure Street, northwest Las Vegas




# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**10<sup>th</sup> Highest** Modeled DMAX8 Day at Joe Neal site, Azure Street, northwest Las Vegas




# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**Highest** Modeled DMAX8 Day at Palo Verde site, Pavilion Center Drive, west Las Vegas




# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**4<sup>th</sup> Highest** Modeled DMAX8 Day at Palo Verde site, Pavilion Center Drive, west Las Vegas



# State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

**10<sup>th</sup> Highest** Modeled DMAX8 Day at Palo Verde site, Pavilion Center Drive, west Las Vegas





# Spatial Distribution of State Ozone Contributions


- Spatial distribution of state's ozone contribution to DMAX8 ozone concentrations greater than or equal to:
  - 76 ppb (current NAAQS)
  - 70 ppb; 65 ppb and 60 ppb (potential future NAAQS)
  - 0 ppb (highest contribution in year)
- Two types of metrics:
  1. Maximum modeled contribution to Highest and 4<sup>th</sup> Highest DMAX8 ozone (from WestJumpAQMS Appendix C)
  2. Attainment Test Unmonitored Areas projection contribution to 8-hour ozone design value (not shown in this presentation)
- Examples for Nevada next:
  - Maximum contribution to highest DMAX8 ever and at 76 ppb (current NAAQS)
  - Maximum contribution to 4<sup>th</sup> high DMAX8 for 76, 70, 65, and 60 ppb

# 2008 Nevada 8-Hour Ozone Contribution

from WestJumpAQMS Appendix C

## Highest Modeled Contribution


Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 0$  ppb  
NV Anthropogenic Max Contribution



Max(68,80) = 27.50

## Modeled DMAX8 Ozone $\geq 76$ ppb

Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 76$  ppb  
NV Anthropogenic Max Contribution




Max(65,83) = 26.29

# 2008 Nevada Contribution to 4<sup>th</sup> High DMAX8 Ozone from WestJumpAQMS Appendix C


**4<sup>th</sup> Highest DMAX8 Ozone  $\geq 76$  ppb**

**4<sup>th</sup> Highest DMAX8 Ozone  $\geq 70$  ppb**

**Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 76$  ppb**  
NV Anthropogenic 4th Highest Contribution



**Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 70$  ppb**  
NV Anthropogenic 4th Highest Contribution




# 2008 Nevada Contribution to 4<sup>th</sup> High DMAX8 Ozone from WestJumpAQMS Appendix C

**4<sup>th</sup> Highest MAX8 Ozone  $\geq 65$ ppb**


**4<sup>th</sup> Highest DMAX8 Ozone  $\geq 60$  ppb**

**Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 65$  ppb**  
NV Anthropogenic 4th Highest Contribution



Max(65,83) = 21.99

**Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 60$  ppb**  
NV Anthropogenic 4th Highest Contribution



Max(65,83) = 21.99


# “Other Sources” Max Contrib. 4<sup>th</sup> High DMAX8 Ozone

Boundary Conditions

Natural


Anthropogenic

Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 0$  ppb




Max(82,2) = 80.37

Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 0$  ppb




Max(70,11) = 12.84

Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 0$  ppb




Max(133,70) = 110.89

Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 0$  ppb




Max(129,53) = 60.13

Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 0$  ppb



Max(116,41) = 6.16

Contrib. to CAMx Daily Max 8-Hour Ozone  $\geq 0$  ppb



Max(79,51) = 3.15


Wildfire

Prescribed Fire


Agricultural Fire

# Northern California Wildfires June-July 2008


June Base08c DMAX8 Ozone Shasta 0007




July Base08c DMAX8 Ozone Shasta 0007



June Base08c DMAX8 Ozone Butte 0007




July Base08c DMAX8 Ozone Butte 0007






# Denver Ozone Monitors July 2008


## Jul DMAX8 Ozone Rocky Flats No




## Jul Base08c DMAX8 Ozone Chatfield



## Jul Base08c DMAX8 Fort Collins West



## Jul Base08c DMAX8 Greeley






### Source Contribution to Annual Average PM<sub>2.5</sub> in ug/m<sup>3</sup>

#### Jarbridge Wilderness


PM<sub>2.5</sub> = 5.79 ug/m<sup>3</sup>; PM<sub>2.5</sub> = 5.79 ug/m<sup>3</sup> (100.0%)



### Composition of Annual Average PM<sub>2.5</sub> in ug/m<sup>3</sup>

#### Jarbridge Wilderness

PM<sub>2.5</sub> = 5.79 ug/m<sup>3</sup>



## Annual Average PM<sub>2.5</sub> (μg/m<sup>3</sup>) Jarbridge Wilderness IMPROVE site

← Sources

Composition


Source category  
Composition example



### Composition of Annual PM<sub>2.5</sub> from Mobile in ug/m<sup>3</sup>

#### Jarbridge Wilderness

PM<sub>2.5</sub> = 5.79 ug/m<sup>3</sup>; Mobile = 0.09 ug/m<sup>3</sup> (1.6%)



# Pilot Study - Detailed Source Category-Specific Ozone Source Apportionment



- Six Source Categories:
  - Natural (Biogenic, Lightning, Sea Salt & WBD)
  - Fires (WF, Rx, & Ag)
  - Upstream Oil and Gas (O&G)
  - Point Sources (EGU & Non-EGU)
  - Mobile Sources (on-road, non-road & CMV)
  - Remainder (Area/Non-Point)
- Ozone Apportionment
  - May-Aug 2008
  - 36/12/4 km Domains
  - 4 States (CO, NM, UT & WY)

# Pilot Study - Ozone Source Category-Specific Source Apportionment

← 4 km Detailed Source Apportionment Domain

36/12/4 km Two-Way Grid Nesting

(Results in Appendix I on WestJumpAQMS webpage)



**CAMx Modeling Domain**

36 km : 148 x 112 (-2736, -2088) to (2592, 1944)  
 12 km\*: 227 x 230 (-2388, -1236) to ( 336, 1524)  
 04 km\*: 164 x 218 (-1228, -436) to ( -572, 436)


\* includes buffer cells

# Detailed Pilot Study: 2008 4<sup>th</sup> Highest Modeled Contribution to Ozone (from WestJumpAQMS Appendix I)


## Canyonlands NP, UT site

### Contributions to MDA8 Ozone [ppb] at UT\_San Juan0101

Rank (4) 05/11/08; Model = 71.2 ppb; Obs = 63.6 ppb; Bias = +12.0%; BC = 64.7 ppb (90.9%)




# WestJumpAQMS Maximum Ozone Season W126 (ppm-hrs)





# Nevada Regional Haze Planning



status of Nevada progress report to EPA ...?

Next regional haze full control SIP due July 2018 (or later)

WestJumpAQMS modeling is the starting point for 2011 or 2014 base year, 2018 progress check, and 2028 projection modeling.

## WestJumpAQMS Benefited From

- WRAP Regional Modeling Center (2002 Platform)
- Four Corners Air Quality Task Force (2005 Platform)
- Continental Divide-Creston EIS (2005/2006 Platform)
  - NEPA O&G EIS using PGM for far-field AQ/AQRV
- Denver Ozone SIP Modeling and Follow-On
- 2008 National Emissions Inventory (2008 NEIv2.0)
  - Cornerstone to 2008 emissions
- WRAP Phase III O&G Emissions Study
  - Projected to 2008 plus add Permian Basin
- WESTAR-funded MEGAN Biogenic Emissions Enhancement Study
- DEASCO<sub>3</sub> 2008 Fire Emissions



## Benefited from WestJumpAQMS

- Colorado Air Resource Management Study (CARMMS)
  - 2008 4 km Modeling Platform
- Deterministic & Empirical Assessment of Smoke's Contribution to Ozone (DEASCO<sub>3</sub>)
  - 2008 36/12 km Modeling Platform
- PMDETAIL -- Smoke contributions to PM
- Three-State Data Warehouse (3SDW) and Three-State Air Quality Study (3SAQS)
  - 2008 36/12/4 km Modeling Platform; Test database for 3SDW
- Additional Follow-On Studies
  - NPS, BLM, etc.

Thanks –

Tom Moore, WRAP Air Quality Program Manager  
Western States Air Resources Council (WESTAR)

e: [tmoore@westar.org](mailto:tmoore@westar.org) | o: 970.491.8837

Western Regional Air Partnership | [www.wrapair2.org](http://www.wrapair2.org)





# Summary of WestJumpAQMS 2008 Modeling Results for Nevada

- Shown earlier
  - Examples of Upwind Ozone Contribution to highest and 4<sup>th</sup> highest, and 10<sup>th</sup> highest modeled days at 5 monitor sites across NV (shown earlier, from Appendix B)
- Next
  - Nitrogen Deposition Analysis
  - Thanks to Tammy M. Thompson, CIRA, and Michael G. Barna, NPS for modeling results and slides

# WestJumpAQMS Modeled Total Nitrogen Deposition

Annual kg N/ha



Class 1 & 2 Areas outlined in **Red**. Total N includes wet & dry deposition of all species.

## Total Modeled Nitrogen Deposition Includes:

- Organic Nitrogen Species: PAN, RNO<sub>3</sub> (model chemistry includes organic N formed from biogenic VOCs)
- Dry Deposition Other Oxidized Nitrogen: NO, NO<sub>2</sub>, N<sub>2</sub>O<sub>5</sub>, HONO, HO<sub>2</sub>NO<sub>2</sub>
- Dry Deposition Ammonia (NH<sub>3</sub>)
- Wet Deposition Other Oxidized Nitrogen: NO, NO<sub>2</sub>, N<sub>2</sub>O<sub>5</sub>, HONO, HO<sub>2</sub>NO<sub>2</sub>
- Dry Deposition of Nitric Acid (HNO<sub>3</sub>)
- Particulate Nitrate (NO<sub>3</sub><sup>-</sup>): Wet and dry
- Particulate Ammonium (NH<sub>4</sub><sup>+</sup>): Wet and dry

## Total Measured Nitrogen Deposition Includes:


- Organic Nitrogen Species: PAN, RNO<sub>3</sub> (model chemistry includes organic N formed from biogenic VOCs)
- Dry Deposition Other Oxidized Nitrogen: NO, NO<sub>2</sub>, N<sub>2</sub>O<sub>5</sub>, HONO, HO<sub>2</sub>NO<sub>2</sub>
- Dry Deposition Ammonia (NH<sub>3</sub>)
- Wet Deposition Other Oxidized Nitrogen: NO, NO<sub>2</sub>, N<sub>2</sub>O<sub>5</sub>, HONO, HO<sub>2</sub>NO<sub>2</sub>
- Dry Deposition of Nitric Acid (HNO<sub>3</sub>)
- Particulate Nitrate (NO<sub>3</sub><sup>-</sup>): Wet and dry
- Particulate Ammonium (NH<sub>4</sub><sup>+</sup>): Wet and dry



# Nitrogen deposition measurement data is incomplete

- Chemical Transport Models (e.g., CAMx) capture the bulk of total Nitrogen Deposition (although CAMx is missing reduced organic nitrogen and includes limited oxidized organic nitrogen – both expected to be small)
- Many Critical Load values are estimated using measurement data, some with scaling factors to account for “missing” N, others incorporate modeling data, still others leave data as is with caveats
- Efforts to close this gap continue, including the creation, in 2010, and subsequent expansion of the Ammonia Monitoring Network (AMoN)

# Critical Load by Eco-Region (kg N/ha)



# Nitrogen Deposition Excess

Total Modeled Nitrogen Wet & Dry Deposition (all species) - Critical Load

