

West-wide Jumpstart Air Quality Modeling Study

Final Project Report and Modeling Results



June 4, 2014

Tom Moore

WRAP Air Quality Program Manager

Technical Project Team

ENVIRON, Alpine Geophysics, Univ. of North Carolina

Funding from State of NM, BP, and National BLM Air Program

Oversight by western states, local air agencies, federal land managers, EPA regional and national offices



EPA national Ozone Standard

- Measured at ground station sites, highest 8-hour average each day
- 4th highest values each year are averaged over 3-year periods to determine compliance (e.g., 2007-09, 2008-10)
 - Statistic is called a “Design Value” for that site for that time period
- Current Ozone health standard level is 75 ppb
- EPA (re)considering revised Ozone health standard in a range of 60 to 70 ppb
- EPA also considering a secondary Ozone standard for ecosystem protection
 - Growing season / daylight hours-weighted cumulative metric

What are (some of) the sources and control issues in the West related to new Ozone standard(s)?


- Urban and rural reactivity
- Transport and formation – how much / how important?
- Public lands with large biogenic emissions and fire activity
 - How to characterize for effects of drought and climate variation ?
- Federal and state mobile fuel and tailpipe controls
- Upstream Gas NSPS rules in place in 2015
 - Industry practices changing rapidly, e.g., green completions
- Point sources (dominated by EGUs for SO₂, NO_x)
 - Significant NO_x BART by ~2018
 - Less coal-fired electricity supply due to climate change rule?
 - 17+ million acres of public lands leased in last 5 years for O&G exploration and production

Introduction

- West-wide Jump-start Air Quality Modeling Study (WestJumpAQMS) was initiated in late 2010 to:
 - Develop the next generation of regional air quality modeling databases for ozone, PM_{2.5}, visibility and deposition planning in the western U.S.
 - Provide information on the role of interstate and international transport to ozone and PM_{2.5} under current and potential future NAAQS
 - Assess contributions of major source categories (e.g., point, O&G, mobile, et cetera) to air quality in the West
 - Provide detailed information to the community

Overview of Approach

- 2008 Modeling Database
 - 36 km CONUS
 - 12 km WESTUS
 - 4 km IMWD
- WRF meteorological; CAMx photochemical; SMOKE emissions models
- 2008 WRAP Phase III O&G emissions
- 2008 NEI emissions
- Model Evaluation
- Sensitivity Tests



- State-Specific and Source Category-Specific Ozone and PM_{2.5} Source Apportionment Modeling

WestJumpAQMS Products

- Final Report
 - 15 Electronic Appendices
 - Response-to-Comments
- Ammonia Emissions Recommendations Memo
- Modeling Protocol
 - Response-to-Comments
- WRF Application/Evaluation Report
 - Evaluation down to individual monitoring site
 - Response-to-Comments
- 16 Technical Memorandums on Emissions
 1. Point Sources
 2. Area + Non-Road
 3. On-Road Mobile
 - 4a-e. Oil and Gas (5 geographic areas)
 5. Fire (WF, Rx & Ag)
 6. Fugitive Dust
 7. Off-Shore Shipping
 8. Ammonia
 9. Biogenic
 - 11 Mexico/Canada
 12. Sea Salt and Lightning
 13. Emissions Modeling Parameters

All information on WestJumpAQMS website

WestJumpAQMS Progress Webinars

- June 20, 2013: 2008 Database Development
- July 26, 2013: State-Specific Source Apportionment
- August 29, 2013: Source Category-Specific Source Apportionment
- Interactive agenda from WestJumpAQMS Final Project Report meeting, Denver, CO – September 25, 2013
- **Presentations and all project materials at:**
<http://www.wrapair2.org/WestJumpAQMS.aspx>

Ozone, PM, Deposition, and Visibility Source Apportionment Resources from WestJumpAQMS

WestJumpAQMS - Reports

West-Wide Jump-Start Air Quality Modeling Study (WestJumpAQMS) – Final Report ([PDF](#) 15MB), September 30, 2013

- Response-to-Comments for Draft Final Report ([PDF](#) 1MB), September 30, 2013

List of Appendices and directions for use ([PDF](#))

Appendix A: CSAPR-Type Analysis for 2008 Upwind State Highest Contribution to Average and Maximum Ozone Design Values at any Monitoring Site in up to 5 Downwind States using MATS ([XLSX](#) 1MB)

Appendix B: State Contributions to Daily Maximum 8-Hour Ozone Concentrations on 10 Highest Modeled Ozone Days in 2008 by Monitoring Site ([XLSX](#) 19MB)

Appendix C: Spatial Maps of State-Specific Anthropogenic Emissions Contributions to Highest and Fourth Highest Modeled Daily Maximum 8-Hour Ozone Concentrations during 2008 Greater than 76 (current NAAQS), 70, 65, 60 (potential future NAAQS) and 0 (maximum contribution) ppb across the 12 km WESTUS and 36 km CONUS Domains ([ZIP](#) 37MB) (**corrected files posted February 7, 2014**)

Appendix D: CSAPR-Type Analysis for 2008 Upwind State Highest Contribution to Annual PM_{2.5} Design Values in up to Five Downwind States using MATS ([XLSX](#) 12MB)

Appendix E: State Contributions to Modeled Annual PM_{2.5} Concentrations in 2008 by Monitoring Site ([XLSX](#) 23MB)

Appendix F: CSAPR-Type Analysis for 2008 Upwind State Highest Contribution to 24-Hour PM_{2.5} Design Values in up to Five Downwind States using MATS ([XLSX](#) 12MB)

Appendix G: Spatial Maps of Modeled State-Specific Anthropogenic Emissions Contributions to Highest (1stmax) and Eighth (8thmax) Highest 24-Hour PM_{2.5} Concentrations during 2008 greater than 35 (current NAAQS), 30, 25, 20 and 0 (maximum contribution) $\mu\text{g}/\text{m}^3$ ([ZIP](#) 13MB) (**corrected files posted February 7, 2014**)

Appendix H: Source Category-Specific Contributions to 8-Hour Ozone Design Values at Monitoring Sites in the 4 km Detailed Source Apportionment Domain (DSAD) using MATS and Maximum Contribution to 8-Hour Ozone Design Values in Each DSAD State (CO, NM, UT and WY) due to Major Source Categories using MATS ([XLSX](#) 1MB)

Appendix I: Source Category-Specific Contributions to Ten Highest Modeled Daily Maximum 8-Hour Ozone Concentrations at Monitoring Sites in the 4 km Detailed Source Apportionment Domain (DSAD) ([XLSX](#) 2MB)

Appendix J: Source Category-Specific Contributions to Annual PM_{2.5} Design Values at Monitoring Sites in the 12 km WESTUS Domain using MATS ([XLSX](#) 2MB)

Appendix K: Source Category-Specific Contributions to Modeled Annual PM_{2.5} Concentrations ($\mu\text{g}/\text{m}^3$) at Monitoring Sites in the 12 km WESTUS Domain ([XLSX](#) 4MB)


Appendix L: Source Category-Specific Contributions to 24-Hour PM_{2.5} Design Values at Monitoring Sites in the 12 km WESTUS Domain using MATS ([XLSX](#) 2MB)

Appendix M: Source Category-Specific Contributions to Ten Highest Modeled 24-Hour PM_{2.5} Concentrations ($\mu\text{g}/\text{m}^3$) at Monitoring Sites in the 12 km WESTUS Domain ([XLSX](#) 10MB)

Appendix N: Annual Sulfur and Nitrogen Wet and Dry Deposition at IMPROVE Monitors by Species ([XLSX](#) 1MB)


Appendix O: Western State-Specific Modeled Contributions to Visibility Impairment at IMPROVE Monitoring Sites for Modeled Worst (W20) and Best (B20) 20% Days during 2008 ([ZIP](#) 46MB)

Ozone Attainment Test Software – Unmonitored Area Analysis with Design Value (2006-2010) ≥ 76 ppb




○ Min(210,3) = 76.00, ◇ Max(45,67) = 113.30


Ozone Attainment Test Software – Unmonitored Area Analysis with Design Value (2006-2010) ≥ 70 ppb



Attainment Test Software – Unmonitored Area Analysis with Design Value (2006-2010) ≥ 65 ppb



Ozone Attainment Test Software – Unmonitored Area Analysis with Design Value (2006-2010) ≥ 60 ppb




State-Specific Ozone Source Apportionment

- Purpose: To provide information on the role of ozone transport to exceedances of the current and potential future ozone NAAQS in the western U.S.
- Approach: Analyze ozone apportionment several ways:
 1. Upwind state contribution to downwind state nonattainment using Cross State Air Pollution Rule (CSAPR-type) approach
 - Use EPA method for projecting ozone Design Values (RRFs)
 2. State contributions to modeled high ozone DMAX8 ozone at monitors in 12 km WESTUS domain
 - Spatial extent of modeled state contributions to 1stmax and 4thmax DMAX8 ozone greater than current and potential future NAAQS
 - Source category analysis (Natural, Fires & Anthropogenic)
 3. Detailed Source Category-Specific Source Apportionment
 - 6 key source categories across 4 states in intermountain West
 - 2-way nesting between model domains

State-Specific Ozone Source Apportionment



- 2008 36/12 km Base
- 17 Western States
 - Plus EasternUS, Can, Mex & Off-Shore
- 5 Source Categories
 - Natural
(Biogenics+Lightning+WBDust+SeaSalt)
 - WF, Rx and Ag Fires
 - Anthropogenic
- 107 Source Groups (21 x 5 + 2)
 - 4 Extra Species for each Group
 - 428 additional species
 - Standard Model = 70 species
 - Computationally Demanding





CSAPR-Type Analysis for current (76 ppb) and potential future (70 and 65 ppb) NAAQS levels

- CSAPR looked at contributions to:
 - Average Design Value = Average of DVs from 2006-2010
 - Max Design Values = Max DVs from 2006-2010
- 136 ozone monitors in 12 km WESTUS domain with Average Design Value exceeding 76 ppb NAAQS
 - 86 sites (63%) in California
- For 17 upwind western states examine 2008 contribution to DMAX8 ozone Design Value in downwind states
 - CSAPR used a 1% NAAQS significance threshold (≥ 0.76 ppb)
- This analysis is for 2008 and is not a regulatory analysis that would have to examine a future year



South Dakota CSAPR-Type Ozone Analysis for potential 70 & 65 ppb NAAQS (from WestJumpAQMS Appendix A)



South Dakota CSAPR-Type Ozone Analysis for potential 70 & 65 ppb NAAQS (from WestJumpAQMS Appendix A)




South Dakota CSAPR-Type Ozone Analysis for potential 65 ppb NAAQS (from WestJumpAQMS Appendix A)




State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

Highest Modeled DMAX8 Day in Blackhawk, SD




State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

4th Highest Modeled DMAX8 Day in Blackhawk, SD



State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

10th Highest Modeled DMAX8 Day in Blackhawk, SD



State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

Highest Modeled DMAX8 Day @ Brookings, SD


Site: SD_Brookings0003

Rank: 1 - 31 May, 2008

Total Ozone = 63.3 ppb

BC Ozone = 39.5 ppb (62.5%)

High Day Contributions to MDA8 Ozone [ppb]



AZ	AZ (0.17 ppb, 0.27%)
CA	CA (0.49 ppb, 0.78%)
CO	CO (0.14 ppb, 0.23%)
KS	KS (0.07 ppb, 0.11%)
ID	ID (0.22 ppb, 0.35%)
MT	MT (0.69 ppb, 1.10%)
OK	OK (0.09 ppb, 0.14%)
OR	OR (0.38 ppb, 0.60%)
WA	WA (0.66 ppb, 1.04%)
WY	WY (0.23 ppb, 0.36%)
ND	ND (2.22 ppb, 3.50%)
SD	SD (0.90 ppb, 1.43%)
NE	NE (0.11 ppb, 0.18%)
NV	NV (0.15 ppb, 0.24%)
UT	UT (0.21 ppb, 0.33%)
TX	TX (0.31 ppb, 0.49%)
NM	NM (0.08 ppb, 0.13%)
Eastern	Eastern (0.72 ppb, 1.14%)
Canada	Canada (8.14 ppb, 12.86%)
Mexico	Mexico (0.16 ppb, 0.25%)
Ocean	Ocean (0.13 ppb, 0.21%)
Natural	Natural (7.26 ppb, 11.48%)
Wild Fire	Wild Fire (0.11 ppb, 0.17%)
Rx Fire	Rx Fire (0.04 ppb, 0.06%)
AG Fire	AG Fire (0.06 ppb, 0.09%)

State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

4th Highest Modeled DMAX8 Day @ Brookings, SD


Site: SD_Brookings0003

Rank: 4 - 03 Apr, 2008

Total Ozone = 60.0 ppb

BC Ozone = 40.6 ppb (67.6%)

High Day Contributions to MDA8 Ozone [ppb]



State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

10th Highest Modeled DMAX8 Day @ Brookings, SD


Site: SD_Brookings0003

Rank: 10 - 21 Aug, 2008

Total Ozone = 58.2 ppb

BC Ozone = 20.0 ppb (34.3%)

High Day Contributions to MDA8 Ozone [ppb]



AZ	AZ (0.06 ppb, 0.11%)
CA	CA (0.07 ppb, 0.12%)
CO	CO (0.08 ppb, 0.14%)
KS	KS (0.23 ppb, 0.40%)
ID	ID (0.03 ppb, 0.06%)
MT	MT (0.05 ppb, 0.08%)
OK	OK (0.11 ppb, 0.19%)
OR	OR (0.02 ppb, 0.04%)
WA	WA (0.03 ppb, 0.05%)
WY	WY (0.06 ppb, 0.10%)
ND	ND (0.10 ppb, 0.18%)
SD	SD (1.59 ppb, 2.74%)
NE	NE (0.26 ppb, 0.45%)
NV	NV (0.03 ppb, 0.05%)
UT	UT (0.07 ppb, 0.12%)
TX	TX (0.46 ppb, 0.78%)
NM	NM (0.10 ppb, 0.17%)
Eastern	Eastern (30.18 ppb, 51.84%)
Canada	Canada (0.31 ppb, 0.54%)
Mexico	Mexico (0.11 ppb, 0.18%)
Ocean	Ocean (0.33 ppb, 0.56%)
Natural	Natural (3.67 ppb, 6.31%)
Wild Fire	Wild Fire (0.08 ppb, 0)
Rx Fire	Rx Fire (0.05 ppb, 0.08%)
AG Fire	AG Fire (0.18 ppb, 0.31%)

State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

Highest Modeled DMAX8 Day at Sioux Falls, SD


Site: SD_Minnehaha0008

Rank: 1 - 31 May, 2008

Total Ozone = 64.1 ppb


BC Ozone = 38.6 ppb (60.2%)

High Day Contributions to MDA8 Ozone [ppb]




State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

4th Highest Modeled DMAX8 Day at Sioux Falls, SD



State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

10th Highest Modeled DMAX8 Day at Sioux Falls, SD



State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

Highest Modeled DMAX8 Day @ Wind Cave National Park, SD


Site: SD_Custer0132

Rank: 1 - 31 May, 2008

Total Ozone = 71.1 ppb

BC Ozone = 51.9 ppb (73.0%)

High Day Contributions to MDA8 Ozone [ppb]



AZ	AZ (0.21 ppb, 0.30%)
CA	CA (3.75 ppb, 5.27%)
CO	CO (0.07 ppb, 0.10%)
KS	KS (0.04 ppb, 0.06%)
ID	ID (0.96 ppb, 1.35%)
MT	MT (0.85 ppb, 1.20%)
OK	OK (0.06 ppb, 0.08%)
OR	OR (0.42 ppb, 0.59%)
WA	WA (0.40 ppb, 0.57%)
WY	WY (2.63 ppb, 3.70%)
ND	ND (0.03 ppb, 0.04%)
SD	SD (0.30 ppb, 0.42%)
NE	NE (0.03 ppb, 0.05%)
NV	NV (0.82 ppb, 1.15%)
UT	UT (0.48 ppb, 0.68%)
TX	TX (0.22 ppb, 0.31%)
NM	NM (0.05 ppb, 0.07%)
Eastern	Eastern (0.27 ppb, 0.38%)
Canada	Canada (0.56 ppb, 0.78%)
Mexico	Mexico (0.18 ppb, 0.26%)
Ocean	Ocean (0.26 ppb, 0.36%)
Natural	Natural (6.45 ppb, 9.07%)
Wild Fire	Wild Fire (0.09 ppb, 0.1)
Rx Fire	Rx Fire (0.06 ppb, 0.08%)
AG Fire	AG Fire (0.01 ppb, 0.01%)

State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

4th Highest Modeled DMAX8 Day @ Wind Cave National Park, SD


Site: SD_Custer0132

Rank: 4 - 01 Jun, 2008

Total Ozone = 68.7 ppb

BC Ozone = 50.4 ppb (73.3%)

High Day Contributions to MDA8 Ozone [ppb]



AZ	AZ (0.19 ppb, 0.27%)
CA	CA (3.22 ppb, 4.69%)
CO	CO (0.07 ppb, 0.10%)
KS	KS (0.04 ppb, 0.06%)
ID	ID (0.88 ppb, 1.28%)
MT	MT (0.86 ppb, 1.25%)
OK	OK (0.06 ppb, 0.08%)
OR	OR (0.47 ppb, 0.68%)
WA	WA (0.45 ppb, 0.65%)
WY	WY (1.98 ppb, 2.89%)
ND	ND (0.03 ppb, 0.04%)
SD	SD (0.88 ppb, 1.28%)
NE	NE (0.04 ppb, 0.06%)
NV	NV (0.71 ppb, 1.03%)
UT	UT (0.44 ppb, 0.64%)
TX	TX (0.21 ppb, 0.31%)
NM	NM (0.05 ppb, 0.07%)
Eastern	Eastern (0.27 ppb, 0.40%)
Canada	Canada (0.67 ppb, 0.98%)
Mexico	Mexico (0.16 ppb, 0.24%)
Ocean	Ocean (0.24 ppb, 0.34%)
Natural	Natural (6.25 ppb, 9.11%)
Wild Fire	Wild Fire (0.08 ppb, 0.1%)
Rx Fire	Rx Fire (0.05 ppb, 0.08%)
AG Fire	AG Fire (0.01 ppb, 0.01%)

State Contributions to Modeled 10 Highest DMAX8 Ozone Days (from WestJumpAQMS Appendix B)

10th Highest Modeled DMAX8 Day @ Wind Cave National Park, SD


Site: SD_Custer0132

Rank: 10 - 16 Apr, 2008

Total Ozone = 65.5 ppb

BC Ozone = 58.3 ppb (89.0%)

High Day Contributions to MDA8 Ozone [ppb]



AZ	AZ (0.00 ppb, 0.00%)
CA	CA (0.26 ppb, 0.40%)
CO	CO (0.00 ppb, 0.00%)
KS	KS (0.00 ppb, 0.00%)
ID	ID (0.17 ppb, 0.26%)
MT	MT (1.44 ppb, 2.20%)
OK	OK (0.00 ppb, 0.00%)
OR	OR (0.31 ppb, 0.47%)
WA	WA (1.14 ppb, 1.74%)
WY	WY (0.34 ppb, 0.51%)
ND	ND (0.00 ppb, 0.00%)
SD	SD (0.11 ppb, 0.17%)
NE	NE (0.00 ppb, 0.00%)
NV	NV (0.02 ppb, 0.03%)
UT	UT (0.00 ppb, 0.01%)
TX	TX (0.00 ppb, 0.00%)
NM	NM (0.00 ppb, 0.00%)
Eastern	Eastern (0.00 ppb, 0.00%)
Canada	Canada (2.03 ppb, 3.10%)
Mexico	Mexico (0.00 ppb, 0.00%)
Ocean	Ocean (0.17 ppb, 0.26%)
Natural	Natural (1.15 ppb, 1.75%)
Wild Fire	Wild Fire (0.00 ppb, 0.00%)
Rx Fire	Rx Fire (0.04 ppb, 0.06%)
AG Fire	AG Fire (0.00 ppb, 0.00%)

Spatial Distribution of State Ozone Contributions

- Spatial distribution of state's ozone contribution to DMAX8 ozone concentrations greater than or equal to:
 - 76 ppb (current NAAQS)
 - 70 ppb; 65 ppb and 60 ppb (potential future NAAQS)
 - 0 ppb (highest contribution in year)
- Two types of metrics:
 1. Maximum modeled contribution to Highest and 4th Highest DMAX8 ozone (from WestJumpAQMS Appendix C)
 2. Attainment Test Unmonitored Areas projection contribution to 8-hour ozone design value (not shown in this presentation)
- Examples for South Dakota next:
 - Maximum contribution to highest DMAX8 ever and at 76 ppb (current NAAQS)
 - Maximum contribution to 4th high DMAX8 for 76, 70, 65, and 60 ppb


2008 South Dakota 8-Hour Ozone Contribution

from WestJumpAQMS Appendix C

Highest Modeled Contribution


DMAX8 Ozone ≥ 76 ppb

Contrib. to CAMx Daily Max 8-Hour Ozone ≥ 0 ppb
SD Anthropogenic Max Contribution



Max(157,146) = 12.72

Contrib. to CAMx Daily Max 8-Hour Ozone ≥ 76 ppb
SD Anthropogenic Max Contribution




Max(159,141) = 1.46

2008 South Dakota Contribution to 4th High DMAX8 Ozone from WestJumpAQMS Appendix C

4th Highest DMAX8 Ozone \geq 76 ppb


4th Highest DMAX8 Ozone \geq 70 ppb

Contrib. to CAMx Daily Max 8-Hour Ozone \geq 76 ppb
SD Anthropogenic 4th Highest Contribution



Max(214,100) = 1.71

Contrib. to CAMx Daily Max 8-Hour Ozone \geq 70 ppb
SD Anthropogenic 4th Highest Contribution




Max(155,144) = 3.06

2008 South Dakota Contribution to 4th High DMAX8 Ozone from WestJumpAQMS Appendix C

4th Highest MAX8 Ozone \geq 65ppb


Contrib. to CAMx Daily Max 8-Hour Ozone \geq 65 ppb
SD Anthropogenic 4th Highest Contribution



Max(157,143) = 6.21

4th Highest DMAX8 Ozone \geq 60 ppb

Contrib. to CAMx Daily Max 8-Hour Ozone \geq 60 ppb
SD Anthropogenic 4th Highest Contribution



Max(201,136) = 7.51


“Other Sources” Max Contrib. 4th High DMAX8 Ozone

Boundary Conditions

Natural


Anthropogenic

Contrib. to CAMx Daily Max 8-Hour Ozone ≥ 0 ppb
BC 4th High Contribution




Max(82,2) = 80.37

Contrib. to CAMx Daily Max 8-Hour Ozone ≥ 0 ppb
Natural 4th High Contribution




Max(70,11) = 12.84

Contrib. to CAMx Daily Max 8-Hour Ozone ≥ 0 ppb
Anthropogenic 4th High Contribution




Max(133,70) = 110.89

Contrib. to CAMx Daily Max 8-Hour Ozone ≥ 0 ppb
Wildfires 4th Highest Contribution




Max(129,53) = 60.13

Contrib. to CAMx Daily Max 8-Hour Ozone ≥ 0 ppb
Rx Burns 4th Highest Contribution



Max(116,41) = 6.16

Contrib. to CAMx Daily Max 8-Hour Ozone ≥ 0 ppb
Agricultural Burns 4th Highest Contribution



Max(79,51) = 3.15


Wildfire

Prescribed Fire


Agricultural Fire

Northern California Wildfires June-July 2008


June Base08c DMAX8 Ozone Shasta 0007




July Base08c DMAX8 Ozone Shasta 0007



June Base08c DMAX8 Ozone Butte 0007




July Base08c DMAX8 Ozone Butte 0007




Denver Ozone Monitors July 2008


Jul DMAX8 Ozone Rocky Flats No




Jul Base08c DMAX8 Ozone Chatfield



Jul Base08c DMAX8 Fort Collins West



Jul Base08c DMAX8 Greeley



Pilot Study - Detailed Source Category-Specific Ozone Source Apportionment



- **Six Source Categories:**
 - Natural (Biogenic, Lightning, Sea Salt & WBD)
 - Fires (WF, Rx, & Ag)
 - Upstream Oil and Gas (O&G)
 - Point Sources (EGU & Non-EGU)
 - Mobile Sources (on-road, non-road & CMV)
 - Remainder (Area/Non-Point)
- **Ozone Apportionment**
 - May-Aug 2008
 - 36/12/4 km Domains
 - 4 States (CO, NM, UT & WY)

Pilot Study - Ozone Source Category-Specific Source Apportionment

← 4 km Detailed Source Apportionment Domain

36/12/4 km Two-Way Grid Nesting

(Results in Appendix I on WestJumpAQMS webpage)



CAMx Modeling Domain

36 km : 148 x 112 (-2736, -2088) to (2592, 1944)
 12 km* : 227 x 230 (-2388, -1236) to (336, 1524)
 04 km* : 164 x 218 (-1228, -436) to (-572, 436)


* includes buffer cells

Detailed Pilot Study: 2008 4th Highest Modeled Contribution to Ozone (from WestJumpAQMS Appendix I)

Canyonlands NP, UT site

Contributions to MDA8 Ozone [ppb] at UT_San Juan0101

Rank (4) 05/11/08; Model = 71.2 ppb; Obs = 63.6 ppb; Bias = +12.0%; BC = 64.7 ppb (90.9%)




Detailed Pilot Study: 2008 4th Highest Modeled Contribution to Ozone (from WestJumpAQMS Appendix I)

Contributions to MDA8 Ozone [ppb] at UT_Uintah1003

Vernal, UT site

Modeled Rank (4) on 06/13/08

no ambient observations, site was not installed



Summary of WestJumpAQMS 2008 Modeling Results for South Dakota

- Shown earlier
 - Examples of Upwind Ozone Contribution to highest and 4th highest, and 10th highest modeled days at 4 monitor sites across SD (shown earlier, from Appendix B)
- Next
 - South Dakota's PM_{2.5} Contributions
 - By Source Category
 - By Species
 - (from Appendix K)


Contributions to Modeled Annual PM2.5 (from WestJumpAQMS Appendix K)

Modeled Annual PM2.5 @ Wind Cave National Park, SD

Source Contribution to Annual Average PM2.5 in ug/m3

SD_Custer0132


PM2.5 = 4.07 ug/m3; PM2.5 = 4.07 ug/m3 (100.0%)



Composition of Annual Average PM2.5 in ug/m3

SD_Custer0132


PM2.5 = 4.07 ug/m3




Contributions to Modeled Annual PM2.5 (from WestJumpAQMS Appendix K)

Modeled Annual PM2.5 @ Badlands National Park, SD

Source Contribution to Annual Average PM2.5 in ug/m3
SD_Jackson0001
PM2.5 = 4.71 ug/m3; PM2.5 = 4.71 ug/m3 (100.0%)



Composition of Annual Average PM2.5 in ug/m3
SD_Jackson0001
PM2.5 = 4.71 ug/m3




Contributions to Modeled Annual PM2.5 (from WestJumpAQMS Appendix K)

Modeled Annual PM2.5 @ Brookings, SD


**Source Contribution to Annual Average PM2.5 in ug/m3
SD_Brookings0002**

PM2.5 = 10.04 ug/m3; PM2.5 = 10.04 ug/m3 (100.0%)



**Composition of Annual Average PM2.5 in ug/m3
SD_Brookings0002**

PM2.5 = 10.04 ug/m3




Contributions to Modeled Annual PM2.5 (from WestJumpAQMS Appendix K)

Modeled Annual PM2.5 @ at Sioux Falls, SD

Source Contribution to Annual Average PM2.5 in ug/m3

SD_Minnehaha0008


PM2.5 = 11.21 ug/m3; PM2.5 = 11.21 ug/m3 (100.0%)



Composition of Annual Average PM2.5 in ug/m3

SD_Minnehaha0008

PM2.5 = 11.21 ug/m3



South Dakota's Regional Haze Program



© 2008 Rikk Flohr

South Dakota's first progress report is due to EPA by January 21, 2016

Next regional haze full control SIP due July 2018

WestJumpAQMS modeling is the starting point for 2011 base year, 2018 progress check, and 2028 projection modeling.

WestJumpAQMS Benefited From

- WRAP Regional Modeling Center (2002 Platform)
- Four Corners Air Quality Task Force (2005 Platform)
- Continental Divide-Creston EIS (2005/2006 Platform)
 - NEPA O&G EIS using PGM for far-field AQ/AQRV
- Denver Ozone SIP Modeling and Follow-On
- 2008 National Emissions Inventory (2008 NEIv2.0)
 - Cornerstone to 2008 emissions
- WRAP Phase III O&G Emissions Study
 - Projected to 2008 plus add Permian Basin
- WESTAR-funded MEGAN Biogenic Emissions Enhancement Study
- DEASCO₃ 2008 Fire Emissions

Benefited from WestJumpAQMS

- Colorado Air Resource Management Study (CARMMS)
 - 2008 4 km Modeling Platform
- Deterministic & Empirical Assessment of Smoke's Contribution to Ozone (DEASCO₃)
 - 2008 36/12 km Modeling Platform
- PMDETAIL -- Smoke contributions to PM
- Three-State Data Warehouse (3SDW) and Three-State Air Quality Study (3SAQS)
 - 2008 36/12/4 km Modeling Platform; Test database for 3SDW
- Additional Follow-On Studies
 - NPS, BLM, etc.

Thanks –

Tom Moore, WRAP Air Quality Program Manager
Western States Air Resources Council (WESTAR)

e: tmoore@westar.org | o: 970.491.8837

Western Regional Air Partnership | www.wrapair2.org